

Odborné články 2008

- Fischer D, Pavlůvčík P, **Sedláček F**, Šálek M, 2009: Predation of the alien American mink, *Mustela vison* on native crayfish in middle-sized streams in central and western Bohemia. Folia Zoologica 58: 45-56.

- Frynta D, Novaková M, Kutalová H, Palme R, **Sedláček F**, 2009: Apparatus for Collection of Fecal Samples from Undisturbed Spiny Mice (*Acomys cahirinus*) Living in a Complex Social Group. J. Am. Assoc. Lab. Anim. Sci. 48: 196-201.

- Knotková E, Veitl S, Šumbera R, **Sedláček F**, Burda H, 2009: Vocalisations of the silvery mole-rat: comparison of vocal repertoires in subterranean rodents with different social systems. Bioacoustics - Int. J. Anim. Sound Record. 18: 241-257.

- Padyšáková E, Šálek M, Poledník L, **Sedláček F**, Albrecht T, 2009: Removal of American mink increases the success of simulated nests in linear habitat. Wildlife Res. 36: 225-230.

2008

- Hulová S, **Sedláček F**, 2008: Population genetic structure of the European ground squirrel in the Czech Republic. Conservation Genetics 9: 615-625.

2007

- Rico A, Kindlmann P, **Sedláček F**, 2007: Barrier effects of roads on movements of small mammals. Folia Zoologica 56: 1-12.

- Rico A, Kindlmann P, **Sedláček F**, 2007: Road crossing in bank voles and yellow-necked mice. Acta Theriologica 52: 85-94.

- Řičánková V, Šumbera R, **Sedláček F**, 2007: Familiarity and partner preferences in female common voles, *Microtus arvalis*. J. Ethology 25: 95-98.

- Zelová J, Šumbera R, **Sedláček F**, Burda H, 2007: Energetics in the solitary subterranean rodent, the silvery mole-rat, *Heliophobius argenteocinereus*, and the allometry of RMR in African mole-rats (*Bathyergidae*). Comp. Biochem. Physiol. A 147: 412-419.

2006

- Zemek F, Heřman M, Kierdorf H, Kierdorf U, **Sedláček F**, 2006: Spatial distribution of dental fluorosis in roe deer (*Capreolus capreolus*) from North Bohemia (Czech Republic) and its relationships with environmental factors. Sci. Total Environment 370: 491-505.

2004

- Hanousková I, Boháč J, **Sedláček F**, Šerá B, Lepšová A, Zacharda M, 2004: Harmful organisms in urban green areas. Ekologia-Bratislava 23, Suppl. 1: 58-68.

2002

- Weiter L, Heřman M, **Sedláček F**, Zemek F, 2002: Potential occurrence of the birch mouse (*Sicista betulina*) in the Bohemian Forest (Šumava): a geographical information system approach. Folia Zoologica 51, Suppl.1: 133-144.

2001

- **Sedláček F**, Zemek F, Heřman M, Kierdorf H, Kierdorf U, 2001: Fluoride Load on Ecosystems in Western Part of Krušné hory mts Determined by Bioindication. Ekologia-Bratislava 20: 70-79.

2000

- Kierdorf H, Kierdorf U, Richards A, **Sedláček F**, 2000: Disturbed Enamel Formation in Wild

Boars (*Sus scrofa L.*) From Fluoride Polluted Areas in Central Europe. The Anatomical Record 259: 12-24.

- Koudela B, Šumbera R, **Sedláček F**, 2000: *Eimeria burdai* sp. n. (Apicomplexa: Eimeriidae), a new parasite species from subterranean African silvery mole-rat, *Heliophobius argenteocinereus*. Folia Parasit. 47: 97-99.

- Zima J, Griffiths H I, Burda H, **Sedláček F**, 2002: Preface - Proceedings of an International Conference Ceske Budejovice, Czech Republic July 10-14, 2000. Folia Zoologica 51, Suppl.1: 2.

1999

- Kierdorf H, Kierdorf U, Sedláček F, 1999: Monitoring regional fluoride pollution in the Saxonian Ore mountains (Germany) using the biomarker dental fluorosis in roe deer (*Capreolus capreolus L.*). The Sci. Total Environ. 232: 159-168.

1998

- Schultz M, Kierdorf U, **Sedláček F**, Kierdorf H, 1998: Pathological bone changes in the mandibles of wild red deer (*Cervus elaphus L.*) exposed to high environmental levels of fluoride. J. Anat. 193: 431-442.

1997

- Kierdorf U, Richards A, **Sedláček F**, Kierdorf H, 1997: Fluoride Content and Mineralization of Red Deer (*Cervus elaphus*) Antlers and Pedicles from Fluoride Polluted and Uncontaminated Regions. Arch. Environ. Contam. Toxicol. 32: 222-227.

1996

- Kierdorf H, Kierdorf U, **Sedláček F**, 1996: Biomonitoring der Fluoridbelastung des Lebensraumes von Wildwiederkauern aus Immissionsgebieten Nordbohmens (Tschechische Republik). Z. Jagdwiss. 42: 41-52.

- Kierdorf U, Kierdorf H, **Sedláček F**, Fejerskov O, 1996: Structural changes in fluorosed dental enamel of red deer (*Cervus elaphus* L.) from a region with severe environmental pollution by fluorides. J. Anat. 188: 183-195.

- Kierdorf H, Kierdorf U, **Sedláček F**, Erdelen M, 1996: Mandibular bone fluoride levels and occurrence of fluoride induced dental lesions in populations of wild red deer (*Cervus elaphus*) from Central Europe. Environmental Pollution 93: 75-81.

1989

- Valach R, **Sedláček F**, 1989: Retention and excretion of fluorides related to their concentration in the drinking-water. In: Anke M et al. (eds.), Proc. 6th Int. Trace Element Symp., K-M. Univ. Leipzig, F-Sch. Univ. Jena 1-5: 1164-1172.

1988

- Boháč J, **Sedláček F**, Švecová Z, Krivoluckij D A, 1988: Animals as bioindicators of industrial emission. Zhurnal obshchei biologii 49: 630-635.

1985

- Molíková D, **Sedláček F**, Bejček V, Šťastný K, Jirouš J, 1985: Changes in the Leucograms of the Common Vole (*Microtus arvalis*) in an Industrially Polluted and in a Background Landscape. Ekológia (ČSFR) 4: 337-345.

Kapitoly v monografiích

2007

- **Sedláček F**, 2007: Adaptive Physiological Mechanisms in the Underground Dwellers. In: Begall S., Burda H., Schleich C.E. (eds), Subterranean Rodents: News from underground. Springer-Verlag, Berlin, Heidelberg, pp. 13-19.

- **Sedláček F**, 2007: New Data on Metabolic Parameters in Subterranean Rodents. In: Begall S., Burda H., Schleich C.E. (eds), Subterranean Rodents: News from underground. Springer-Verlag, Berlin, Heidelberg, pp. 35-47.

2005

- Tomášková L, Bejček V, **Sedláček F**, Šťastný K, Tkadlec E, Zima J, 2005: Population biology of shrews (*Sorex araneus* and *S. minutus*) from a polluted area in central Europe. In: Merritt J.F. et al. (eds): Advances in the Biology of Shrews II. Int. Soc. Shrew Biologists, pp. 189-197.

1994

- **Sedláček F**, Bejček V, 1994: An Approach to Biomonitoring the Industrial Pollution of the Landscape Using the Common Vole (*Microtus arvalis*). In: Biological Monitoring of the Environment: A Manual of Methods. CAB Int. IUBS, Wallingford, pp. 87-90.

1989

- Šťastný K, Bejček V, Paukert J, **Sedláček F**, 1989: Bioindikation – Einsatz- und Aussagemöglichkeiten bei der Umweltüberwachung. In: Geoökologische Stoffflüsse und Konsequenzen anthropogener Aktivitäten in der Landschaft. VEB Deutscher Verlag für Grundstoffindustrie, Leipzig, pp. 110-123.